Comment on "$K^-d \to \pi\Sigma n$ reactions and structure of the $\Lambda(1405)$" by S. Ohnishi, Y. Ikeda, T. Hyodo, E. Hiyama, W. Weise

J. Révai[1] and N.V. Shevchenko[2]

[1] *Wigner Research Center for Physics, RMI, H-1525 Budapest, P.O.B. 49, Hungary*
[2] *Nuclear Physics Institute, 25068 Řež, Czech Republic*

(Dated: August 5, 2014)

PACS numbers:

arXiv:1408.0661v1 [nucl-th] 4 Aug 2014

We are astonished by the main statement of the paper [1]:
"We report on the first results of a full three-body calculation of the $\bar{K}NN \to \pi YN$ amplitude for the $K^-d \to \pi\Sigma n$ reaction, and examine how the $\Lambda(1405)$ resonance manifests itself in the neutron energy distributions of the $K^-d \to \pi\Sigma n$ reactions."

A calculation of the same reaction within the same theoretical framework and with the same goal was reported at several international conferences and published in [2] without being mentioned in [1].

The $\bar{K}N$ potentials applied in the two calculations differ from each other, the quality of their reproduction of experimental data favors those of [2]. In addition, a doubtful approximation was used in the potentials of some of the authors of [1] in their previous papers, which, as it seems, were used in [1] too. A discussion on this issue can be found in [3].

The original parts of [1] in comparison with [2] are higher partial waves and an energy-dependent potential. From this point of view [1] seems to be a natural extension of the work done in [2]. However the use of nonrelativistic AGS equations for initial kaon momentum 1 GeV/c is highly questionable. Therefore, the realization of the author's promise to take relativistic corrections into account should have preceded the publication.

---

[1] S. Ohnishi, Y. Ikeda, T. Hyodo, E. Hiyama, W. Weise, "$Kd \to \pi\Sigma n$ reactions and structure of the $\Lambda(1405)$", arXiv:1408.0118 (2014)

[2] J. Révai, "Signature of the $\Lambda(1405)$ resonance in neutron spectra from the $K^- + d$ reaction", Few-Body Syst. **54** (2013) 1865

[3] J. Révai, N.V. Shevchenko, "Faddeev calculations of the $\bar{K}NN$ system with chirally-motivated $\bar{K}N$ interaction. II. The Kpp quasi-bound state", arXiv:1403.0757 (2014)